

Information following:

Endodontic Treatment

Root Canals

- Endodontic treatment can take 1, 2, or 3 appointments depending on the case. It is possible to experience any of the following symptoms after any of these appointments: sensitivity to hot and/or cold; sensitivity to pressure; possible swelling. If you experience swelling, please call our office immediately as additional medication may be required.
- One common occurrence with a newly rootcanaled tooth is for the tooth to feel high when you bite your teeth together. If this occurs it will cause your tooth to stay sensitive for a longer period of time. Please call us if your bite feels "high" as this problem is easily solved with a simple bite adjustment.

(337) 837-1861

300 W Madison St – Broussard, LA 70518 Matthew Melancon, D.D.S. Melancondentistry.com

- A temporary filling may be used to temporarily seal the tooth between visits.
- Be gentle on the tooth while eating until the final restoration is placed.
- The gum tissue could have been irritated during the procedure and may be sore for a few days together with the anesthetic injection site.
- During endodontic treatment, the nerve, blood and nutrient supply to the tooth is removed. This will cause the tooth to become brittle and prone to fracturing which could result in the need to extract the tooth. A full coverage crown is recommended to prevent this from happening.
- Proper brushing, flossing, and regular 6-month (minimal) cleanings are essential to the longterm stability of your teeth.